

The mathfont Package in Action: Two Mathematical Snippets Rendered in Times New Roman  
Conrad Kosowsky

Mathematicians usually define  $e$  in one of two ways: as the horizontal asymptote of a certain function or as the limit of an infinite series. Specifically, it's most common to see  $e$  defined as either

$$e = \lim_{x \rightarrow \infty} \left(1 + \frac{1}{x}\right)^x$$

or

$$e = \sum_{k=0}^{\infty} \frac{1}{k!}.$$

The first definition is simpler in that it involves a limit of a single expression, not a limit of partial sums, but in practice, the second tends to be more tractable. The power series expression of  $e^x$  is given by

$$\sum_{n=0}^{\infty} \frac{x^n}{n!},$$

and the relationship between this expression and the series definition is much more apparent than it is for the first limit. This relationship arises in a variety of different mathematical contexts, for example the famous Euler's formula  $e^{i\theta} = \cos \theta + i \sin \theta$  or the related definition of the characteristic function for a random variable  $X$ :

$$\phi_X(t) = \mathbb{E} \left( e^{iX} \right).$$

Expanding  $e^{iX}$  as a power series gives an expression for  $\phi_X$  that we can differentiate term by term.

A smooth manifold consists of a topological space  $M$  equipped with a smooth maximal atlas  $\{\phi_i\}$ . The maps  $\phi_i: U_i \rightarrow \mathbb{R}^n$  are continuous, and their compositions  $\phi_i \circ \phi_j^{-1}$  are diffeomorphisms on subsets of  $\mathbb{R}^n$ . If we have a map  $f: M \rightarrow N$  between manifolds, this structure allows us to talk about differentiability of  $f$ . Specifically, we say that  $f$  is smooth if for any  $i$  and  $j$ , the composition

$$\psi_j \circ f \circ \phi_i^{-1}$$

is itself smooth, where  $\{\psi_j\}$  is a smooth atlas for  $N$ . Differentiating  $f$  produces the associated tangent map  $Df$ . The function  $Df$  maps the tangent bundle  $TM$  to the tangent bundle  $TN$  and is linear when restricted to individual tangent spaces  $T_pM$ . If  $M$  can be written as a product  $M_1 \times M_2$ , we can consider the partial tangent maps  $\partial_1 f$  and  $\partial_2 f$  by considering the compositions  $f \circ \iota_1$  and  $f \circ \iota_2$ , where  $\iota_1$  and  $\iota_2$  are inclusion maps with respect to a particular point. Combining both maps, we have the equation

$$Df(u, v) = \partial_1 f(u) + \partial_2 f(v),$$

and this relationship can be thought of as an adaption of the standard product rule.